Table 19

Reasons for Treatment by Age of Patients - DI in 2018

(Based on the information on Data Collection Forms received from licensed centre up to 31 December 2018)

| Reasons | Age Group (by no. of patients) | | | | | | | | | |
|---|--------------------------------|-------|-------|-------|-------|-------|-------|-------|--------------|-----|
| | 18-20 | 21-25 | 26-30 | 31-35 | 36-40 | 41-45 | 46-50 | 51-55 | 56 and above | All |
| Obstructive azoospermia | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Nonobstructive azoospermia | 0 | 0 | 0 | 3 | 0 | 0 | 0 | 0 | 0 | 3 |
| Severe deficits in semen quality in couples who do not wish to undergo intracytoplasmic sperm injection | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Genetic | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Infectious disease in the male partner (such as HIV) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Severe rhesus isoimmunisation | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Others | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

Remarks:

- (1) The age of husband has been used in calculating the age of patient.
- (2) Same patient might have more than one reason for treatment.